

Owner's Copy

PennHIP Report

Referring Veterinarian: Dr Todd Baker

Email: ftmacvet@hotmail.com

Clinic Name: Fort Macleod Veterinary Clinic

Clinic Address: 609 12 St

Fort Macleod, AB TOL 0Z0

Phone: (403) 553-4887 Fax: (403) 553-2924

Patient Information

Client: Payne, Kelsie

Patient Name: Dreamlands Mr Darcy Reg. Name: Dreamlands Mr Darcy

PennHIP Num: 120813 Species: Canine

Date of Birth: 02 Jan 2018

Sex: Male

Date of Study: 03 Aug 2018

Date of Report: 06 Aug 2018

Tattoo Num:

Patient ID: Dreamlands Mr Darcy

Registration Num:

Microchip Num: 900026000636532

Breed: LABRADOODLE

Age: 7 months

Weight: 20.2 lbs/9.2 kgs Date Submitted: 03 Aug 2018

Findings

Distraction Index (DI): Right DI = 0.49, Left DI = 0.48.

Osteoarthritis (OA): No radiographic evidence of OA for either hip.

Cavitation/Other Findings:None.


Interpretation

Distraction Index (DI): The laxity ranking is based on the hip with the greater laxity (larger DI). In this case the DI used is

OA Risk Category: The DI is between 0.31 and 0.49. This patient is at mild risk for hip OA.

Distraction Index Chart:

LABRADOODLE



Distraction Index

Breed Statistics: This interpretation is based on a cross-section of 6422 canine patients of the LABRADOODLE breed in the AIS PennHIP database. The gray strip represents the central 90% range of DIs (0.32 - 0.76) for the breed. The breed average DI is 0.51 (solid square). The patient DI is the solid circle (0.49).

Summary: The degree of laxity (DI = 0.49) falls within the central 90% range of DIs for the breed. This amount of hip laxity places the hip at a mild risk to develop hip OA. No radiographic evidence of OA for either hip.

Interpretation and Recommendations: No OA/Mild Risk: Low risk to develop radiographic evidence of hip OA early in life, however OA may manifest after 6 years of age or later. Risk of OA increases as DI, age, body weight, and activity level increase. OA susceptibility is breed specific, larger breeds being more susceptible. Recommendations: Evidencebased strategies to lower the risk of dogs developing hip OA or to treat those having OA fall into 5 modalities.* For